

FINDINGS & RECOMMENDATIONS FROM THE INTER-ASSOCIATION TASK FORCE FOR APPROPRIATE CARE OF THE SPINE

Mission of the Summit:

To develop guidelines for the pre-hospital management of the physically active with suspected spinal injury.

GENERAL GUIDELINES

- * Any athlete suspected of having a spinal injury should not be moved and should be managed as though a spinal injury exists.
- * The athlete's airway, breathing and circulation, neurological status and level of consciousness should be assessed.
- * The athlete should not be moved unless absolutely essential to maintain airway, breathing, and circulation.
- * If the athlete must be moved to maintain airway, breathing and circulation, the athlete should be placed in a supine position while maintaining spinal immobilization.
- * When moving a suspected spine injured athlete, the head and trunk should be moved as a unit. One accepted technique is to manually splint the head to the trunk.
- * The Emergency Medical Services system should be activated.

FACE MASK REMOVAL

- * The face mask should be removed prior to transportation, regardless of current respiratory status.
- * Those involved in the pre-hospital care of injured football players should have the tools for face mask removal readily available.

FOOTBALL HELMET REMOVAL

The athletic helmet and chin strap should only be removed ...

- * if the helmet and chin strap do not hold the head securely, such that immobilization of the helmet does not also immobilize the head.
- * if the design of the helmet and chin strap is such that even after removal of the face mask the airway can not be controlled, or ventilation be provided.
- * if the face mask can not be removed after a reasonable period of time.
- * if the helmet prevents immobilization for transportation in an appropriate position.

HELMET REMOVAL

Spinal immobilization must be maintained while removing the helmet.

- * Helmet removal should be frequently practiced under proper supervision.
- * Specific guidelines for helmet removal need to be developed.
- * In most circumstances, it may be helpful to remove the cheek padding and/or deflate air padding prior to helmet removal.

EQUIPMENT

Appropriate spinal alignment must be maintained.

- * There needs to be a realization that the helmet and shoulder pads elevate an athlete's trunk when in the supine position.
- * Should either be **removed**, or if only one is present, appropriate spinal alignment must be maintained.
- * The front of the shoulder pads can be opened to allow access for CPR and defibrillation.

This task force encourages the development of a local emergency care plan regarding the pre-hospital care of the athlete with suspected spine injury. This plan should include communication with the institution's administration and those directly involved with the assessment and transportation of the injured athlete.

All providers of pre-hospital care should practice and be competent in all the skills identified in these guidelines before they are needed in an emergency situation.

These guidelines were developed as a consensus statement by;

Douglas M. Kleiner, PhD, ATC, FACSM (Chair), National Athletic Trainers' Association; **Jon Almquist, ATC**, National Athletic Trainers' Association Secondary School Athletic Trainers Committee; **Julian Bailes, MD**, American Association of Neurological Surgeons; **John C. Biery, DO, FAOSM**, American Osteopathic Academy of Sports Medicine; **Pepper Burruss, ATC, PT**, Professional Football Athletic Trainers' Society; **Alexander M. Butman, DSc, REMT-P**, National Registry of Emergency Medical Technicians; **Jerry Diehl**, National Federation of State High School Associations; **Robert Domeler, MD**, National Association of Emergency Medical Services Physicians; **Kent Falb, ATC, PT**, National Athletic Trainers' Association; **Henry Feuer, MD**, National Football League Physicians Society; **Jay Greenstein, DC**, American Chiropractic Association Council on Sports Injuries and Physical Fitness; **Letha Y. Griffin, MD**, American Orthopaedic Society for Sports Medicine; National Collegiate Athletic Association Committee on Competitive Safeguards and Medical Aspects of Sports; **Bob Hannemann, MD**, American Academy of Pediatrics Committee on Sports Medicine and Fitness; **Margaret Hunt, ATC**, United States Olympic Committee; **Daniel Kraft, MD**, American Medical Society for Sports Medicine; **James Laughnane, ATC**, National Athletic Trainers' Association College and University Athletic Trainers' Committee; **Connie McAdam, MICT**, National Association of Emergency Medical Technicians; **Dennis A. Miller, ATC, PT**, National Athletic Trainers' Association; **Michael Oliver**, National Operating Committee on Safety and Equipment; **Andrew N. Pollack, MD**, OrthopaedicTrauma Association; **Dan Smith, DPT, ATC**, American Physical Therapy Association Sports Physical Therapy Section; **David Thorson, MD**, American Academy of Family Physicians; **Patrick R. Trainor, ATC**, National Association of Intercollegiate Athletics; **Robert G. Watkins, MD**, American Academy of Orthopaedic Surgeons Committee on the Spine; **Stuart Weinstein, MD**, American College of Sports Medicine; North American Spine Society; Physiatrix Association of Spine, Sports, and Occupational Rehabilitation.

June 04, 1998

Please disregard all previous drafts.